Web Images Videos Maps News Shopping Gmail more ▼	Scholar Preferences Sign in
COOSIC SCHOOL	
Scholar Articles and patents - 2003 include citations Create email alert	Results 1 - 10 of about 90. (0.08 sec)
Topical ALA-photodynamic therapy for the treatment of acne vulgaris W Hongchan, CR Taylor, Y Chang Journal of Investigative, 2000 - nature.com Bacterial porphyrin fluorescence was also suppressed by photodynamic therapy gland function, and apparent decrease in follicular bacteria after photodynamic therapy ALA, aminolevulinic acid; PDT, photodynamic therapy; PpIX, protoporphyrin IX; SO, sebum output Cated by 253 - Related articles - Bt. Difect - All 5 versions	
Photodynamic therapy of acne vulgaris with topical δ -aminolaevulinic acid and incoherent light in	
Japanese patients Y Hon, Y Nihomya, B Tajima. — British Journal of …, 2001 – Wiley Online Library Repeated 5-aminolevulinic acid-based photodynamic therapy for pigmented basal cell carcinoma. J Dermatol 2000; 27: 10 – 15 R. In vivo photoproduct formation during PDT with ALA-induced endogenous porphyrins . J Photochem Photobiol B 1993; 18: 287 – 90 Cited by 123 – Britanes autoles – 8L Orest – ALC versions	
Acne phototherapy with a high-intensity, enhanced, narrow-band, blue light source; an open study and in vitro investigation A Kawada, Y Aragare. H Kameyame Journal of, 2002 - Elsevier Topical ALA-photodynamic therapy is effective for acne vulgaris with significant side effects such as transient hyperpigmentation, superficial exfoliation, and Blue light therapy for acne may be a kind of photodynamic therapy using endogenous porphyrins produced by P Sitest by 81 - Bulesed studies - Alt 21 sersions	[PBF] from verilux.com
Topical 5-aminolaevulinic acid photodynamic therapy for the treatment of skin conditions other than non-melanoma skin cancer SH libbotson - British Journal of Dematology, 2002 - Wiley Online Library permeability of abnormal stratum corneum, relative iron deficiency and alterations in porphyrin enzyme profiles in NMSC, non-melanoma skin cancers; ALA, 5-aminolaevulinic acid; PDT , photodynamic therapy Acne vulgaris, Keratoacanthoma, Epidermodysplasia verruciformis Catact by 78 - Beisted Judicies - 88, Girset - Alt Z. Netschas	IEBELfran. sansitosia.com
Eradication of Propionibacterium acnes by its endogenic porphyrins after illumination with high intensity blue light H Ashkenazi, Z Mailik, Y Harth FEMS Immunology &, 2003 - Wiley Online Library vulgaris with visible light. Dermatology 194, 256–260 [34] Zeina, B., Greenman, J., Purcell, WM, Das, B. (2001) Killing of cutaneous microbial species by photodynamic therapy. Br. J. Dermatol. 144, 274–278 [35] Ramstad, S., Futsaether, CM, Johnsson, A. (1997) Porphyrin Grant by 134 - Related articles - All 12 versions	IPDEL from Jubris.com
Treatment of cutaneous leishmaniasis with photodynamic therapy CD Enk, C Fritsch, F Jones Archives of, 2003 - Am Med Assoc recently demonstrated in a randomized double-blind trial of ALA-PDT for the 9 The use of porphyrins in combination with the quinonoid compound menadione causes Photodynamic therapy is an attractive antiparasitic therapeutic modality that offers rapid localized destruction Cited by 38 - Rewind articles - IN, fared - Alia versions	
A pilot study of ICG laser therapy of acne vulgaris; photodynamic and photothermolysis treatment VV Tuonin, BA Genina, AN Bashkatov Lasers in surgery 2003 - Witey Online Library Photodynamic therapy (PDT) is a well-known procedure used for recovery of many abnormalities including tumors The main toxic agents produced by PDT are singlet oxygen or other reactive acnes colonies taken from the skin site at excitation of bacteria produced porphyrins Cited by 58 - Periated studies - St. Direct - Alt 2 versions.	IPDFI Irom palomat ru
The study of endogenous porphyrins in human skin and their potential for photodynamic therapy by laser induced fluorescence spectroscopy. K Köng, H Meyer, H Someckenburger Lasers in Medical Science, 1993 - Springer protoporphyrin. It seems to be possible to use the endogenous porphyrins for a photodynamic therapy of acne vulgaris and to monitor the therapeutic effect by the simultaneous measurement of spectral changes. INTRODUCTION Cited by 5 - Polated studies - St. Direct - All & versions	
Photodynamic therapy and topical aminolevulinic acid: an overview AK Gunta American Journal of Clinical, 2003 - Ingentaconnect.com AK = actinic keratosis; PDT = photo- dynamic therapy. © Adis Data Information BV 2003 produce endogenous porphyrins (eg coproporphyrin III).[45] Coproporphyrin III can absorb visible blue light at 415nm, sug of aminolevulinic acid/photodynamic therapy in acne vulgaris: Chied by 33 - Related articles - Bt. Cirect - At 3 versions	IPBELfram. adisoniine.com
Apparatus and method for high energy photodynamic therapy of acne vulgaris and seborrhea Y Harth US Patent App. 09/756,130, 2001 - Google Patents [0003] The enlargement and obstruction of sebaceous glands cause acne vulgaris 410 nanometer range), or less efficiently, red (630-670 nanom- eter range) are able to induce a photodynamic effect in which the porphyrins in the [0005] Photodynamic therapy (PDT) is based	

Ched by 11 - Related spicies - All 4 versions

Create email alert

G000000008 € №
Result Page: 1 2 3 4 5 6 7 8 9 Next

porphyrin PDT OR "photodynamic the Search

Go to Google Home - About Google - About Google Scholar

©2011 Google